



A Study on $b_{i,j}^*$ -Open Set within Ideal Biopological Spaces

Ibtissam R. Bukhatwa *

Department of Mathematics, Faculty of Education, University of Benghazi, Benghazi, Libya

دراسة حول المجموعات $b_{i,j}^*$ -المفتوحة داخل الفضاءات ثنائية التبولوجيا المثلية

إبتسام رجب بوخطوة
قسم الرياضيات، كلية التربية، جامعة بنغازي، بنغازي، ليبيا

*Corresponding author: ibtissam.bukhatwa@uob.edu.ly

Received: September 25, 2025 | Accepted: December 07, 2025 | Published: December 20, 2025

Abstract:

This paper introduces the concepts of $b_{i,j}^*$ -I-open sets and $b_{i,j}^*$ -I-closed sets in the framework of ideal ditopological spaces, presenting them as a new form of generalized open and closed sets. $b_{i,j}^*$ -I-open sets are defined as subsets satisfying specific generalized openness conditions with respect to two topologies and an ideal, while $b_{i,j}^*$ -I-closed sets are their complements. The study begins with a precise definition of these sets and explains how they extend previously developed ideas in generalized and bitopological structures. It examines their relationships with other known types of open and closed sets and highlights essential properties that characterize their behavior. Particular attention is given to how the interaction of the two topologies and the ideal shapes their structure and flexibility. The paper also introduces the related operators, $b_{i,j}^*$ -I-interior and $b_{i,j}^*$ -I-closure, and discusses their key properties. Overall, the study contributes to understanding openness and closedness in ideal bitopological spaces and provides a foundation for further research.

Keywords: Ideal ditopological spaces, b -I-open sets, Interior operators, Closure operators, Generalized open sets.

الملخص

تقدم هذه الورقة مفاهيم المجموعات $b_{i,j}^*$ -المفتوحة و $b_{i,j}^*$ -المغلقة في إطار الفضاءات ثنائية الطوبولوجيا مع وجود مثالي، باعتبارها شكلًا جديداً من المجموعات المفتوحة والمغلقة في الطوبولوجيا المعممة. تعرف المجموعات المفتوحة $b_{i,j}^*$ -b على أنها المجموعات التي تحقق شروط الانفتاح المعمم بالنسبة لطوبولوجيين اثنين والمثالي، بينما تعرف المجموعات المغلقة على أنها المكملات الخاصة بها. تبدأ الدراسة بتعريف دقيق لهذه المجموعات وشرح كيفية توسيعها للأفكار المطورة سابقاً في البنى الطوبولوجية الثنائية والمعممة. كما تبحث الدراسة علاقاتها بالمجموعات المفتوحة والمغلقة المعروفة وتبرز الخصائص الأساسية التي تحدد سلوكها. ويُولى اهتمام خاص لتأثير تفاعل الطوبولوجيين والمثالي على بنية هذه المجموعات ومرورتها ضمن الفضاء.

وبعد ذلك تقدم الدراسة أيضا العاملين المرتبطين بهذه المجموعات، وهما $b_{i,j}^* \mathcal{I} - \text{interior}$ و- $\mathcal{I} - b_{i,j}^*$ closure، وتناقش الخصائص الأساسية المرتبطة بهما. تساهم الدراسة في تعزيز الفهم العميق لانفتاح والانغلاق في الفضاءات ثنائية الطوبولوجيا المثالية وتتوفر قاعدة للبحوث المستقبلية.

الكلمات المفتاحية: الفضاءات ثنائية التبولوجيا المثالية، المجموعات المفتوحة، معامل الداخل، معامل الانغلاق، المجموعات المفتوحة المعممة.

Introduction

Kelly introduced the concept (Kelly, 1963) of a ditopological space $(\Xi, \mathcal{T}_1, \mathcal{T}_2)$, where a set Ξ is equipped with two topologies \mathcal{T}_1 and \mathcal{T}_2 . In 1966 (Kuratowski, 2014) studied the notion of ideals on a topological space (Ξ, \mathcal{T}) , defined as collections of subsets of Ξ with the hereditary (i) if $\mathcal{A} \in \mathcal{I}$ and $\mathcal{B} \subset \mathcal{A}$, then $\mathcal{B} \in \mathcal{I}$ and (ii) if $\mathcal{A} \in \mathcal{I}$ and $\mathcal{B} \in \mathcal{I}$ then $\mathcal{A} \cup \mathcal{B} \in \mathcal{I}$. Let $(\Xi, \mathcal{T}_1, \mathcal{T}_2, \mathcal{I})$ an ideal bitopological space consider the power set $\mathcal{P}(\Xi)$ which consists of all subsets of Ξ .

Janković[12], define a set operator $(.)_i^* : \mathcal{P}(\Xi) \rightarrow \mathcal{P}(\Xi)$ called the local function of \mathcal{A} with respect to \mathcal{T}_i and \mathcal{I} , as for $\mathcal{A} \subset \Xi$, $\mathcal{A}_i^*(\mathcal{T}_i, \mathcal{I}) = \{x \in \Xi \mid \mathcal{B} \cap \mathcal{A} \notin \mathcal{I}, \text{ for every } \mathcal{B} \in \mathcal{T}_i(x)\}$ as well as $\mathcal{T}_i(x) = \{\mathcal{B} \in \mathcal{T}_i \mid x \in \mathcal{B}\}$. For each ideal topological space $(\Xi, \mathcal{T}_1, \mathcal{T}_2, \mathcal{I})$, there exists a topology \mathcal{T}_i^* that is finer than \mathcal{T}_i , generated by the base by the base $\mathbb{B}(\mathcal{I}, \mathcal{T}) = \{\mathcal{A} \setminus I \mid \mathcal{A} \in \mathcal{T} \text{ and } I \in \mathcal{I}\}$, However, in general, $\mathbb{B}(\mathcal{I}, \mathcal{T})$ does not always form a topology.

Furthermore, observe that the closure operator corresponding to the topology $\mathcal{T}_i^*(\mathcal{I})$, which is finer than \mathcal{T}_i , is defined by $\text{Cl}_i^*(\mathcal{A}) = \mathcal{A} \cup \mathcal{A}_i^*$.

Accordingly, $\text{Int}_i^*(\mathcal{A})$ denotes the interior of \mathcal{A} in the topology $\mathcal{T}_i^*(\mathcal{I})$ while, $\text{Int}_i^*(\mathcal{A}_i^*)$ represents the interior of \mathcal{A}_i^* with respect to the topology \mathcal{T}_i^* .

The study of open sets in ideal and ditopological spaces has evolved through several key contributions. In 2007, Akdag (Akdag, 2007) introduced the concept of $\mathfrak{b}\text{-}\mathcal{I}$ -open sets" in ideal topological spaces, while in the same year, Al-Hawary and Al-Omari (Al-Hawary & Al-Omari, 2007) proposed the notion of $\mathfrak{b}_{i,j}$ -open sets in bitopological spaces. Building on these ideas, in (Sarma, 2015) expanded this framework by defining $\mathfrak{b}_{i,j}\mathcal{I}$ -open sets in ideal bitopological spaces. Several researchers have also discussed the concept of open sets, as presented in the studies listed in references (Abdullah & Mohammed, 2020), (Ashraf, 2025), (Bukhatwa & Demiralp, 2024), and (Sarma & Acharjee, 2021). These works have contributed significantly to the development of the fundamental concepts and the expansion of the theoretical framework within ditopological spaces.

Preliminaries

In this paper, $(\Xi, \mathcal{T}_1, \mathcal{T}_2, \mathcal{I})$ constantly mean bitopological space with no separation axioms are assumed in this space and TB-S be the abbreviation for ideal bitopological space. And assume \mathcal{A} is a subset of Ξ , then we denote

respectively the interior and the closure of \mathcal{A} by $\text{int}_i(\mathcal{A})$ and $\text{Cl}_j(\mathcal{A})$ with reference to \mathbb{V}_i for $i = 1, 2$.

Definition 2.1. (Al-Hawary & Al-Omari, 2007) Assume \mathcal{A} is a subset of the bitopological space $(\Xi, \mathbb{T}_1, \mathbb{T}_2)$ such that $i, j = 1, 2$, and $i \neq j$. Then \mathcal{A} is called:

$b_{i,j}$ -OS if $\mathcal{A} \subseteq \text{Int}_i(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i(\mathcal{A}))$. (Abdullah & Mohammed, 2020).

Definition 2.2. Assume \mathcal{A} is a subset of the TB-S such that $i, j = 1, 2$, and $i \neq j$. Then \mathcal{A} is called:

1- $\mathbb{I}_{i,j}$ -OS if $\mathcal{A} \subseteq \text{Int}_i(\mathcal{A}_j^*)$. (Caldas et al., 2018).

2- $\text{semi}_{i,j}\mathbb{I}$ -OS if $\mathcal{A} \subseteq \text{Cl}_j^*(\text{Int}_i(\mathcal{A}))$ (Caldas et al., 2020).

3- $\text{semi}_{i,j}^*\mathbb{I}$ -OS if $\mathcal{A} \subseteq \text{Cl}_j(\text{Int}_i^*(\mathcal{A}))$ (Bukhatwa & Demiralp, 2020)

4- $\text{pre}_{i,j}\mathbb{I}$ -OS if $\mathcal{A} \subseteq \text{Int}_i(\text{Cl}_j^*(\mathcal{A}))$. (Caldas et al., 2011)

5- $b_{i,j}\mathbb{I}$ -OS if $\mathcal{A} \subseteq \text{Int}_i(\text{Cl}_j^*(\mathcal{A})) \cup \text{Cl}_j^*(\text{Int}_i(\mathcal{A}))$. (Sarma, 2015)

6- $\alpha_{i,j}\mathbb{I}$ -OS if $\mathcal{A} \subseteq \text{Int}_i(\text{Cl}_j^*(\text{Int}_i(\mathcal{A})))$. (El-Maghrabi et al., 2017)

Lemma 2.1.[8] Let $(\Xi, \mathbb{T}_1, \mathbb{T}_2, \mathbb{I})$ be a TB-S, and $\mathcal{A} \subset \Xi$, then for any $(i, j)\mathbb{I}$ -open set we get

1- $\text{Cl}_j^*(\mathcal{A}) \subset \text{Cl}_j(\mathcal{A})$.

2- $\text{Int}_i(\mathcal{A}) \subset \text{Int}_i^*(\mathcal{A})$.

Lemma 2.2.[8] If $(\Xi, \mathbb{T}_1, \mathbb{T}_2, \mathbb{I})$ be a TB-S, and $\mathcal{A} \subset \Xi$, with reference to any $(i, j)\mathbb{I}$ -OS then we get $\mathcal{A}_j^* = (\text{int}_i(\mathcal{A}_j^*))_j^*$.

Lemma 2.3.[10] Assume \mathcal{A} is a subset of the TB-S $(\Xi, \mathbb{T}_1, \mathbb{T}_2, \mathbb{I})$, and $\mathcal{B} \in \mathbb{T}_j$. Then, $\text{Cl}_j(\mathcal{A}) \cap \mathcal{B} \subseteq \text{Cl}_j(\mathcal{A} \cap \mathcal{B})$.

3- ON $b_{i,j}^*\mathbb{I}$ -OPEN SET

Definition 3.1. If $(\Xi, \mathbb{T}_1, \mathbb{T}_2, \mathbb{I})$ is an ideal bitopological spaces then a subset \mathcal{A} of X is called:

$b_{i,j}^*\mathbb{I}$ -OS if $\mathcal{A} \subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A}))$,

in which $i, j \in \{1, 2\}$, $i \neq j$. The collection comprised of all $b_{i,j}^*\mathbb{I}$ -OS in Ξ will be denoted by $b_{i,j}^*\mathbb{I}$ -O(Ξ), and TB-S for ideal bitopological spaces.

Example 3.1. Consider TB-S such that ; $\Xi = \{v, \mu, \omega, \tau\}$ and $\mathbb{T}_1 = \{\phi, \{\mu\}, \{v, \omega, \tau\}, \Xi\}$, $\mathbb{T}_2 = \{\phi, \{v, \mu\}, \Xi\}$. And if $\mathbb{I} = \{\phi, \{\omega\}\}$. Then $\{v, \omega\}$ and $\{v, \mu, \tau\}$ are $b_{1,2}^*\mathbb{I}$ -OS, but $\{\omega, \tau\}$ is not $b_{1,2}^*\mathbb{I}$ -OS.

Example 3.2. Let IB-S such that ; $\Xi = \{v, \mu, \omega\}$ and $T_1 = \{\phi, \{v\}, \{\mu\}, \{v, \mu\}, \Xi\}$, $T_2 = \{\phi, \{v, \omega\}, \Xi\}$. And if $\mathcal{I} = \{\phi, \{\mu\}, \{\omega\}, \{\mu, \omega\}\}$. Then $\{\mu, \omega\}$ is $b_{1,2}^* \mathcal{I}$ -OS, but $\{\mu, \omega\}$ is not $b_{2,1}^* \mathcal{I}$ -OS.

Example 3.3. Let IB-S such that ; $\Xi = \{v, \mu, \omega, \tau\}$ and $T_1 = \{\phi, \{\mu\}, \{v, \omega\}, \{v, \mu, \omega\}, \Xi\}$, $T_2 = \{\phi, \{\omega\}, \Xi\}$. And if $\mathcal{I} = \{\phi, \{v\}, \{\mu\}, \{v, \mu\}\}$. Then $\{v, \mu, \omega\}$ and $\{v, \tau\}$ are a $b_{1,2}^* \mathcal{I}$ -OS, but $\{\tau\}$ is not $b_{1,2}^* \mathcal{I}$ -OS.

Example 3.4. Let IB-S such that ; $\Xi = \{v, \mu, \omega\}$ and $T_1 = \{\phi, \{\omega\}, \Xi\}$, $T_2 = \{\phi, \{v\}, \{v, \omega\}, \Xi\}$. And if $\mathcal{I} = \{\phi, \{\mu\}, \{\omega\}, \{\mu, \omega\}\}$. Then all subset of Ξ are $b_{2,1}^* \mathcal{I}$ -OS.

Proposition3.1. If $(\Xi, T_1, T_2, \mathcal{I})$ is a IB-S , and \mathcal{A} a subset of Ξ . Then the following properties are observed:

- 1- Every $\mathcal{I}_{i,j}$ -OS is $b_{i,j}^* \mathcal{I}$ -OS.
- 2- Every $b_{i,j} \mathcal{I}$ -OS is $b_{i,j}^* \mathcal{I}$ -OS.
- 3- Every $\text{semi}_{i,j} \mathcal{I}$ -OS is $b_{i,j}^* \mathcal{I}$ -OS.
- 4- Every $\text{pre}_{i,j} \mathcal{I}$ -OS is $b_{i,j}^* \mathcal{I}$ -OS.
- 5- Every $\alpha_{i,j} \mathcal{I}$ -OS is $b_{i,j}^* \mathcal{I}$ -OS.
- 6- Every $\text{semi}_{i,j}^* \mathcal{I}$ -OS is $b_{i,j}^* \mathcal{I}$ -OS.

Proof. (1) Assume \mathcal{A} is a $\mathcal{I}_{i,j}$ -OS, so $\mathcal{A} \subset \text{Int}_i(\mathcal{A}_j^*) \subset \text{Int}_i(\mathcal{A}_j^* \cup \mathcal{A}) \subset \text{Int}_i(\text{Cl}_j^*(\mathcal{A})) \subset \text{Int}_i^*(\text{Cl}_j^*(\mathcal{A})) \subset \text{Int}_i^*(\text{Cl}_j(\mathcal{A}))$, so $\mathcal{A} \subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A}))$.

The remaining proofs are follow from the first statement.

Let $(\Xi, T_1, T_2, \mathcal{I})$ be a IB-S . Assume \mathcal{A} is a subset of Ξ then we get this diagram:




Figure 1: The figure illustrates the relationships between the open sets in the ideal ditopological spaces.

Generally, the opposites of proposition 3.1 is not accurate. as demonstrated by the next example.

Example 3.5. From example 3.1 we have $\{v, \omega\}$ is $b_{1,2}^* \mathcal{I}$ -OS, but it is not $\text{semi}_{i,j} \mathcal{I}$ -OS.

Example 3.6. From example 3.2 we have $\{\mu, \omega\}$ is $b_{1,2}^* I$ -OS, but it is not $b_{i,j} I$ -OS.

Theorem 3.1. If (Ξ, T_1, T_2, I) is a TB-S, let \mathcal{A} be a subset of Ξ . If \mathcal{A} is $b_{i,j}^* I$ -OS with $\text{Int}_i^*(\text{Cl}_j(\mathcal{A})) = \emptyset$, then \mathcal{A} is semi $b_{i,j}^* I$ -OS.

Proof. Let \mathcal{A} be $b_{i,j}^* I$ -OS then $\mathcal{A} \subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A})) \subseteq \text{Cl}_j(\text{Int}_i^*(\mathcal{A}))$.

Theorem 3.2. If (Ξ, T_1, T_2, I) is a TB-S, let \mathcal{A} , and \mathcal{B} be a subsets of X . If \mathcal{A} is $b_{i,j}^* I$ -OS and is a pre $b_{i,j}^* I$ -OS, then $\mathcal{A} \cup \mathcal{B}$ is $b_{i,j}^* I$ -OS.

Proof.

$$\begin{aligned}
 \mathcal{A} \cup \mathcal{B} &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A})) \cup \text{Int}_i(\text{Cl}_i^*(\mathcal{B})) \\
 &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A})) \cup \text{Int}_i^*(\text{Cl}_j(\mathcal{B})) \\
 &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A}) \cup \text{Cl}_j(\mathcal{B})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A})) \cup \\
 &\quad \text{Int}_i^*(\mathcal{B}) \\
 &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A} \cup \mathcal{B})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A} \cup \mathcal{B})).
 \end{aligned}$$

So. $\mathcal{A} \cup \mathcal{B}$ is $b_{i,j}^* I$ -OS.

Theorem 3.3. If (Ξ, T_1, T_2, I) is a TB-S, let \mathcal{A} , and \mathcal{B} be a subsets of X . If \mathcal{A} is pre $b_{i,j}^* I$ -OS and \mathcal{B} is semi $b_{i,j}^* I$ -OS, then $\mathcal{A} \cup \mathcal{B}$ is $b_{i,j}^* I$ -OS.

Proof.

$$\begin{aligned}
 \mathcal{A} \cup \mathcal{B} &\subseteq \text{Int}_i(\text{Cl}_j^*(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{B})) \\
 &\subseteq \text{Int}_i(\text{Cl}_j^*(\mathcal{A} \cup \mathcal{B})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A}) \cup \\
 &\quad \text{Int}_i^*(\mathcal{B})) \\
 &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A} \cup \mathcal{B})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A} \cup \mathcal{B})).
 \end{aligned}$$

So. $\mathcal{A} \cup \mathcal{B}$ is $b_{i,j}^* I$ -OS.

Theorem 3.4. If (Ξ, T_1, T_2, I) is a TB-S, let \mathcal{A} , and \mathcal{B} be a subsets of X . If \mathcal{A} is $b_{i,j}^* I$ -OS and $\mathcal{B} \in T_i$, then $\mathcal{A} \cap \mathcal{B}$ is $b_{i,j}^* I$ -OS.

Proof. Let \mathcal{A} be $b_{i,j}^* I$ -OS, so $\mathcal{A} \subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A}))$

$$\begin{aligned}
 \text{Then } \mathcal{A} \cap \mathcal{B} &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A})) \cap \mathcal{B} \\
 &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A}) \cap \mathcal{B}) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A}) \cap \mathcal{B}) \\
 &\subseteq \text{Int}_i^*(\text{Cl}_j(\mathcal{A} \cap \mathcal{B})) \cup \text{Cl}_j(\text{Int}_i^*(\mathcal{A} \cap \mathcal{B})).
 \end{aligned}$$

Theorem 3.5. If (Ξ, T_1, T_2, I) is a TB-S. Then the union of any $b_{i,j}^* I$ -OS is a $b_{i,j}^* I$ -OS.

Proof. Let $\mathcal{A}_\alpha \in b_{i,j}^* I$ -OS(Ξ) for each $\alpha \in \Lambda$, where Λ is an index set

$$\begin{aligned}
\mathcal{A}_\alpha &\subseteq \text{Int}_i^* \left(\text{Cl}_j(\mathcal{A}_\alpha) \right) \cup \text{Cl}_j \left(\text{Int}_i^*(\mathcal{A}_\alpha) \right), \text{ thus} \\
\bigcup_{\alpha \in \Lambda} \mathcal{A}_\alpha &\subseteq \bigcup_{\alpha \in \Lambda} \left\{ \text{Int}_i^* \left(\text{Cl}_j(\mathcal{A}_\alpha) \right) \cup \text{Cl}_j \left(\text{Int}_i^*(\mathcal{A}_\alpha) \right) \right\} \\
&\subseteq \left\{ \text{Int}_i^* \left(\bigcup_{\alpha \in \Lambda} \text{Cl}_j(\mathcal{A}_\alpha) \right) \right\} \quad \cup \\
\left(\text{Cl}_j \left(\bigcup_{\alpha \in \Lambda} \text{Int}_i^*(\mathcal{A}_\alpha) \right) \right) &\subseteq \left\{ \text{Int}_i^* \left(\text{Cl}_j \left(\bigcup_{\alpha \in \Lambda} \mathcal{A}_\alpha \right) \right) \right\} \quad \cup \\
\text{Cl}_j \left(\text{Int}_i^* \left(\bigcup_{\alpha \in \Lambda} \mathcal{A}_\alpha \right) \right) &.
\end{aligned}$$

Then $\bigcup_{\alpha \in \Lambda} \mathcal{A}_\alpha$ is $b_{i,j}^* \mathcal{I}$ -OS.

Definition 3.2. If $(\Xi, T_1, T_2, \mathcal{I})$ is a TB-S, let \mathcal{A} be a subset of X . \mathcal{A} is called a $b_{i,j}^* \mathcal{I}$ -closed set if its complement is a $b_{i,j}^* \mathcal{I}$ -OS. All $b_{i,j}^* \mathcal{I}$ -CS in Ξ will be denoted by $b_{i,j}^* \mathcal{I}C(\Xi)$.

Theorem 3.6. If $(\Xi, T_1, T_2, \mathcal{I})$ is a TB-S, $\mathcal{A} \subset \Xi$, then \mathcal{A} is a $b_{i,j}^* \mathcal{I}$ -CS if,

$$\text{Cl}_i^*(\text{Int}_j(\mathcal{A})) \cap \text{Int}_j(\text{Cl}_i^*(\mathcal{A})) \subseteq \mathcal{A}.$$

Proof. Assume \mathcal{A} is a $b_{i,j}^* \mathcal{I}$ -CS of $(\Xi, T_1, T_2, \mathcal{I})$, then $(\Xi - \mathcal{A})$ is $b_{i,j}^* \mathcal{I}$ -OS and since

$$\begin{aligned}
\Xi - \mathcal{A} &\subseteq \text{Int}_i \left(\text{Cl}_j^*(\Xi - \mathcal{A}) \right) \cup \text{Cl}_j^* \left(\text{Int}_i(\Xi - \mathcal{A}) \right) \\
&\subseteq \text{Int}_i \left(\text{Cl}_j(\Xi - \mathcal{A}) \right) \cup \text{Cl}_j \left(\text{Int}_i(\Xi - \mathcal{A}) \right) \\
&\subseteq \{\Xi - \text{Cl}_i^*(\text{Int}_j(\mathcal{A}))\} \cup \{\Xi - (\text{Int}_i(\text{Cl}_j^*(\mathcal{A})))\} \\
&\subseteq \Xi - \{\text{Cl}_i^*(\text{Int}_j(\mathcal{A})) \cap \text{Int}_j(\text{Cl}_i^*(\mathcal{A}))\}, \text{ so}
\end{aligned}$$

$$\text{Cl}_i^*(\text{Int}_j(\mathcal{A})) \cap \text{Int}_j(\text{Cl}_i^*(\mathcal{A})) \subseteq \mathcal{A}.$$

Theorem 3.7. If $(\Xi, T_1, T_2, \mathcal{I})$ is a TB-S, let \mathcal{A} and \mathcal{B} be subsets of X . If \mathcal{A} and \mathcal{B} are $b_{i,j}^* \mathcal{I}$ -CS, then $\mathcal{A} \cap \mathcal{B}$ is $b_{i,j}^* \mathcal{I}$ -CS.

Proof. Let \mathcal{A} and \mathcal{B} be $b_{i,j}^* \mathcal{I}$ -CS, so

$$\begin{aligned}
\{\text{Cl}_i^*(\text{Int}_j(\mathcal{A})) \cap \text{Int}_j(\text{Cl}_i^*(\mathcal{A}))\} \cap \{\text{Cl}_i^*(\text{Int}_j(\mathcal{B})) \cap \text{Int}_j(\text{Cl}_i^*(\mathcal{B}))\} &\subseteq \mathcal{A} \cap \mathcal{B} \\
\{\text{Cl}_i^*(\text{Int}_j(\mathcal{A})) \cap \text{Cl}_i^*(\text{Int}_j(\mathcal{B}))\} \cap \{\text{Int}_j(\text{Cl}_i^*(\mathcal{A})) \cap \text{Int}_j(\text{Cl}_i^*(\mathcal{B}))\} &\subseteq \mathcal{A} \cap \mathcal{B} \\
\text{Cl}_i^*(\text{Int}_j(\mathcal{A} \cap \mathcal{B})) \cap \text{Int}_j(\text{Cl}_i^*(\mathcal{A} \cap \mathcal{B})) &\subseteq \mathcal{A} \cap \mathcal{B}
\end{aligned}$$

Definition 3.3. If $(\Xi, T_1, T_2, \mathcal{I})$ is a TB-S, let \mathcal{A} be a subset of X , and x be a point of X . Then,

1- x is called an $b_{i,j}^* \mathcal{I}$ -interior point of \mathcal{A} if there exists $\mathcal{B} \in b_{i,j}^* \mathcal{I}$ -OS such that $x \in \mathcal{B} \subset \mathcal{A}$.

2- The set of all $b_{i,j}^* \mathcal{I}$ -interior points of \mathcal{A} is called $b_{i,j}^* \mathcal{I}$ -interior of \mathcal{A} and is denoted by $b_{i,j}^* \mathcal{I}\text{-Int}(\mathcal{A})$.

Theorem 3.8. Let (Ξ, T_1, T_2, I) be a TB-S, and let \mathcal{A} and \mathcal{B} be subsets of (Ξ, T_1, T_2, I) . The following properties are observed.

- 1- $b_{i,j}^* I\text{-Int}(\mathcal{A}) = \cup \{U: U \subset \mathcal{A} \text{ and } \mathcal{A} \in b_{i,j}^* I\text{-O}(\Xi)\}$.
- 2- $b_{i,j}^* I\text{-Int}(\mathcal{A})$ is the largest $b_{i,j}^* I\text{-OS}$ subset of Ξ contained in \mathcal{A} .
- 3- \mathcal{A} is $b_{i,j}^* I\text{-OS}$ if and only if $\mathcal{A} = b_{i,j}^* I\text{-Int}(\mathcal{A})$.
- 4- $b_{i,j}^* I\text{-Int}(b_{i,j}^* I\text{-Int}(\mathcal{A})) = b_{i,j}^* I\text{-Int}(\mathcal{A})$.
- 5- If $\mathcal{A} \subset \mathcal{B}$, then $b_{i,j}^* I\text{-Int}(\mathcal{A}) \subset b_{i,j}^* I\text{-Int}(\mathcal{B})$.
- 6- $b_{i,j}^* I\text{-Int}(\mathcal{A} \cap \mathcal{B}) \subset b_{i,j}^* I\text{-Int}(\mathcal{A}) \cap b_{i,j}^* I\text{-Int}(\mathcal{B})$.
- 7- $b_{i,j}^* I\text{-Int}(\mathcal{A}) \cup b_{i,j}^* I\text{-Int}(\mathcal{B}) \supset b_{i,j}^* I\text{-Int}(\mathcal{A}) \cup b_{i,j}^* I\text{-Int}(\mathcal{B})$.

Proof. (1) Let $x \in \cup \{U: U \subset \mathcal{A} \text{ and } \mathcal{A} \in b_{i,j}^* I\text{-O}(\Xi)\}$, then there exist $U \subset b_{i,j}^* I\text{-O}(\Xi)$, such that $x \in U \subset \mathcal{A}$. Since $x \in b_{i,j}^* I\text{-Int}(\mathcal{A})$. We get $x \in \cup \{U: U \subset \mathcal{A} \text{ and } \mathcal{A} \in b_{i,j}^* I\text{-O}(\Xi)\} \subset b_{i,j}^* I\text{-Int}(\mathcal{A})$. In the reverse direction, the argument proceeds as follows.

$x \in b_{i,j}^* I\text{-Int}(\mathcal{A})$, then there exist $U \subset b_{i,j}^* I\text{-O}(\Xi)$, such that $x \in U \subset \mathcal{A}$. Then $\in \cup \{U: U \subset \mathcal{A} \text{ and } \mathcal{A} \in b_{i,j}^* I\text{-O}(\Xi)\}$. So $b_{i,j}^* I\text{-Int}(\mathcal{A}) \subset \cup \{U: U \subset \mathcal{A} \text{ and } \mathcal{A} \in b_{i,j}^* I\text{-O}(\Xi)\}$, then we get $b_{i,j}^* I\text{-Int}(\mathcal{A}) = \cup \{U: U \subset \mathcal{A} \text{ and } \mathcal{A} \in b_{i,j}^* I\text{-O}(\Xi)\}$,

(6) Since $\mathcal{A} \cap \mathcal{B} \subset \mathcal{A}$ and $\mathcal{A} \cap \mathcal{B} \subset \mathcal{B}$, by (4), we have $b_{i,j}^* I\text{-Int}(\mathcal{A} \cap \mathcal{B}) \subset b_{i,j}^* I\text{-Int}(\mathcal{A})$ and $b_{i,j}^* I\text{-Int}(\mathcal{A} \cap \mathcal{B}) \subset b_{i,j}^* I\text{-Int}(\mathcal{B})$.

So $b_{i,j}^* I\text{-Int}(\mathcal{A} \cap \mathcal{B}) \subset b_{i,j}^* I\text{-Int}(\mathcal{A}) \cap b_{i,j}^* I\text{-Int}(\mathcal{B})$.

(7). We have $b_{i,j}^* I\text{-Int}(\mathcal{A}) \subset b_{i,j}^* I\text{-Int}(\mathcal{A} \cup \mathcal{B})$ and $b_{i,j}^* I\text{-Int}(\mathcal{B}) \subset b_{i,j}^* I\text{-Int}(\mathcal{A} \cup \mathcal{B})$, then, we have $b_{i,j}^* I\text{-Int}(\mathcal{A}) \cup b_{i,j}^* I\text{-Int}(\mathcal{B}) \subset b_{i,j}^* I\text{-Int}(\mathcal{A} \cup \mathcal{B})$.

The remaining proofs can be established directly.

Definition 3.4. Let (Ξ, T_1, T_2, I) be a TB-S, and $\mathcal{A} \subset \Xi$, and x be a point of Ξ . Then,

- 1- x is called an $b_{i,j}^* I$ -cluster point of \mathcal{A} if $\mathcal{A} \cap B \neq \emptyset$ for every $B \in b_{i,j}^* I\text{-OS}(\Xi, x)$.
- 2- The set of all $b_{i,j}^* I$ -cluster point of \mathcal{A} is called $b_{i,j}^* I$ -cluster of \mathcal{A} and is denoted by $b_{i,j}^* I\text{-Cl}(\mathcal{A})$.

Theorem 3.9. Let (Ξ, T_1, T_2, I) be a TB-S, and let \mathcal{A} and \mathcal{B} be subsets of (Ξ, T_1, T_2, I) . The following properties are observed:

- 1- $b_{i,j}^* I\text{-Cl}(\mathcal{A}) = \cap \{U: \mathcal{A} \subset U \text{ and } U \in b_{i,j}^* I\text{-C}(\Xi)\}$.
- 2- $b_{i,j}^* I\text{-Cl}(\mathcal{A})$ is the smallest $b_{i,j}^* I$ -closed subset of Ξ containing \mathcal{A} .
- 3- \mathcal{A} is $b_{i,j}^* I$ -closed if and only if $\mathcal{A} = b_{i,j}^* I\text{-Cl}(\mathcal{A})$.
- 4- $b_{i,j}^* I\text{-Cl}(b_{i,j}^* I\text{-Cl}(\mathcal{A})) = b_{i,j}^* I\text{-Cl}(\mathcal{A})$.
- 5- If $\mathcal{A} \subset \mathcal{B}$ then $b_{i,j}^* I\text{-Cl}(\mathcal{A}) \subset b_{i,j}^* I\text{-Cl}(\mathcal{B})$.

6- $b_{i,j}^* I\text{-Cl}(\mathcal{A} \cap \mathcal{B}) \subset b_{i,j}^* I\text{-Cl}(\mathcal{A}) \cap b_{i,j}^* I\text{-Cl}(\mathcal{B})$.

7- $b_{i,j}^* I\text{-Cl}(\mathcal{A} \cup \mathcal{B}) \supset b_{i,j}^* I\text{-Cl}(\mathcal{A}) \cup b_{i,j}^* I\text{-Cl}(\mathcal{B})$.

Proof. The proofs can be established directly.

Theorem 3.10. Let (Ξ, T_1, T_2, I) be a IB -S, and let \mathcal{A} be subsets of (Ξ, T_1, T_2, I) , the following properties are observed:

1- $b_{i,j}^* I\text{-Int}(\Xi - \mathcal{A}) = \Xi - b_{i,j}^* I\text{-Cl}(\mathcal{A})$.

2- $b_{i,j}^* I\text{-Cl}(\Xi - \mathcal{A}) = \Xi - b_{i,j}^* I\text{-Int}(\mathcal{A})$.

Proof. (1) Let $x \in b_{i,j}^* I\text{-Cl}(\mathcal{A})$, then there exist $\mathcal{B} \subset b_{i,j}^* I\text{-OS}(\Xi)$, such that $\mathcal{A} \cap \mathcal{B} = \emptyset$. We get $x \in b_{i,j}^* I\text{-Int}(\Xi - \mathcal{A})$, so $\Xi - b_{i,j}^* I\text{-Cl}(\mathcal{A}) \subset b_{i,j}^* I\text{-Int}(\Xi - \mathcal{A})$. Let $x \in b_{i,j}^* I\text{-Int}(\Xi - \mathcal{A})$, we know $b_{i,j}^* I\text{-Int}(\Xi - \mathcal{A}) \cap \mathcal{B} = \emptyset$, $x \notin b_{i,j}^* I\text{-Cl}(\mathcal{A})$, then $x \in \Xi - b_{i,j}^* I\text{-Cl}(\mathcal{A})$. Therefore $b_{i,j}^* I\text{-Int}(\Xi - \mathcal{A}) = \Xi - b_{i,j}^* I\text{-Cl}(\mathcal{A})$. (2). This result can be deduced from (1).

Conclusion

In conclusion, this study has systematically developed the theory of $b_{(i,j)}^* I$ -open and $b_{(i,j)}^* I$ -closed sets, thereby establishing a new and coherent framework for analysis within ideal ditopological spaces. The concomitant formulation of the $b_{(i,j)}^* I$ -interior and $b_{(i,j)}^* I$ -closure operators has deepened the structural understanding of generalized open sets in this dual-topology setting, successfully extending several key results. Ultimately, these contributions do not merely generalize existing concepts but provide a vital and precise foundation for future investigations into advanced topological properties, including novel forms of continuity, compactness, and separation axioms specifically tailored for ideal ditopological systems.

References

1. Abdullah, B. S., & Mohammed, A. A. (2020). *ii-Open set in bitopological spaces*. Rafidain Journal of Computer and Mathematics, 14(2), 33.
2. Akdag, M. (2007). On b-I-open sets and b-I-continuous functions. *International Journal of Mathematics and Mathematical Sciences*, 2007, Article ID 75721. <https://doi.org/10.1155/2007/75721>.
3. Al-Hawary, T., & Al-Omari, A. (2007). b-open and b-continuity in bitopological spaces. *Al-Manarah*, 13(3), 89–101.
4. Ashraf, M. M. (2025). Applications of pre-open & α -open sets in bitopological spaces. *Journal of Mathematics and Statistics Studies*, 6(3), 40-49.
5. Bukhatwa, I., & Demiralp, S. (2020). On some generalized open sets in ideal bitopological spaces. *Journal of New Theory*, (32), 1–10.
6. Bukhatwa, I., & Demiralp, S. (2024). On Strong $\$(i, j)$ $\$$ -Semi $\$^{\{*\}}$ $\$$ - $\$ \backslash \Gamma$ $\$$ -Open Sets in Ideal Bitopological Space. *Journal of New Theory*, (46), 89-98.

7. Caldas, M., Jafari, S., & Rajesh, N. (2011). Preopen sets in ideal bitopological spaces. *Bulletin of Parana's Mathematical Society*, 29(2), 61-68.
8. Caldas, M., Jafari, S., & Rajesh, N. (2020). Semi-open sets in ideal bitopological spaces. *An. Univ. Sci. Budapest.*, (to appear).
9. Caldas, M., Jafari, S., Rajesh, N., & Smarandache, F. (2018). On I-OS and I-continuous functions in ideal bitopological spaces. In *Collected Papers, Volume XIV: Neutrosophics and Other Topics* (p. 338).
10. Dvalishvili, B. (2005). *Bitopological spaces: Theory, relations with generalized algebraic structures and applications* (Vol. 199). Elsevier.
11. El-Maghrabi, A. I., Caldas, M., Jafari, S., Latif, R. M., Nasef, A., Rajesh, N., & Shanthi, S. (2017). Properties of ideal bitopological α -open sets. *Scientific Studies and Research. Series Mathematics and Informatics*, 27(2), 15–36.
12. Janković, D., & Hamlett, T. R. (1990). New topologies from old via ideals. *The American Mathematical Monthly*, 97(4), 295–310.
13. Kelly, J. C. (1963). Bitopological spaces. *Proceedings of the London Mathematical Society*, s3-13(1), 71–89.
14. Kuratowski, K. (2014). *Topology: Volume II*. Elsevier.
15. Sarma, D. J. (2015). bI-open sets in ideal bitopological spaces. *International Journal of Pure and Applied Mathematics*, 105(1), 7–18.
16. Sarma, D. J., & Acharjee, S. (2021). Weakly semi-I-open sets in ideal bitopological spaces. *Transactions of the National Academy of Sciences of Azerbaijan, Series: Physical-Technical and Mathematical Sciences - Mathematics*, 41(1), 71–77.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of **SJPHRT** and/or the editor(s). **SJPHRT** and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.